CORRECTION

Correction: The purine nucleoside phosphorylase *pnp-1* regulates epithelial cell resistance to infection in *C. elegans*

Eillen Tecle, Crystal B. Chhan, Latisha Franklin, Ryan S. Underwood, Wendy Hanna-Rose, Emily R. Troemel

After publication of this article, the authors noted an error in the amino acid change predicted for the *pnp-1(jy90)* mutant allele. The mutant allele is incorrectly annotated as Leucine throughout the text. The wild-type codon is TCC, which codes for the amino acid Serine, and the correct *jy90* mutant allele is TTC, which codes for the amino acid Phenylalanine. The *PLOS Pathogens* editors have confirmed that this error does not affect the conclusions of the study.

Fig 1 is incorrect due to the error described above. The authors have provided a corrected version here.



G OPEN ACCESS

Citation: Tecle E, Chhan CB, Franklin L, Underwood RS, Hanna-Rose W, Troemel ER (2022) Correction: The purine nucleoside phosphorylase *pnp-1* regulates epithelial cell resistance to infection in *C. elegans.* PLoS Pathog 18(7): e1010699. https://doi.org/10.1371/journal.ppat.1010699

Published: July 7, 2022

Copyright: © 2022 Tecle et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Fig 1. pnp-1 mutants have increased expression of IPR genes. A) Gene structure of the two isoforms of pnp-1 with exons indicated as black boxes. 5' and 3' untranslated regions are not shown. B-D) pals-5p:gfp IPR reporter expression in wild-type animals, pnp-1(jy90) and pnp-1(jy121) mutants. myo-2p::mCherry is a pharyngeal marker for the presence of the IPR reporter transgene. Scale bar is $100~\mu m$. E) qRT-PCR of a subset of IPR genes in pnp-1(jy90) and pnp-1 (jy121) mutants. Fold change in gene expression is shown relative to wild-type animals. Graph shows the mean fold change of three independent experiments. Error bars are standard deviation (SD). Mixed stage populations of animals were used. **** indicates p < 0.0001 by one-tailed t-test. F) Quantification of inosine and hypoxanthine levels in pnp-1 and pals-22 mutants from metabolomics analysis. Graph shows the mean levels of metabolites from six independent experiments for pnp-1(jy121), pnp-1(jy90) and pals-22(jy1) mutants, and five independent experiments for wild-type animals. Error bars are standard error of the mean (SEM). ** indicates p < 0.01 by the Kruskal-Wallis test. E, F) Red dots indicate values from individual experiments. See materials and methods for more information.

https://doi.org/10.1371/journal.ppat.1010699.g001

In the *pnp-1* is a negative regulator of IPR gene expression subsection of the Results, there is an error in the fourth sentence of the first paragraph. The correct sentence is: From this analysis, we identified a missense mutation in the PNP gene *pnp-1*, which should result in substitution of a conserved serine (S51 or S68 in isoform a or b, respectively) to phenylalanine.

There is an error in the sixth sentence of the fifth paragraph of the Discussion. The correct sentence is: Support for the model that the catalytic activity of pnp-1 is required for its effects on the IPR comes from the pnp-1(jy90) allele, which has a conserved serine mutated to phenylalanine.

In the Forward mutagenesis screening and cloning of pnp-1(jy90) subsection of the Methods, there is an error in the last sentence of the first paragraph. The correct sentence is: pnp-1(jy90) contains a G to A substitution that should convert serine 51 to phenylalanine in isoform A and serine 68 to phenylalanine in isoform B.

Reference

 Tecle E, Chhan CB, Franklin L, Underwood RS, Hanna-Rose W, Troemel ER (2021) The purine nucleoside phosphorylase *pnp-1* regulates epithelial cell resistance to infection in *C. elegans*. PLoS Pathog 17(4): e1009350. https://doi.org/10.1371/journal.ppat.1009350 PMID: 33878133