S8 Table. Association of MSM and HET with transmission

clusters. The percentages of individuals found to be associated with transmission clusters in MSM and HET in several studies are collated. In the second column, the numbers in parantheses indicate sample sizes examined. Where available, P values and the largest cluster sizes are indicated as other details.

Region	Group (n)	Associated with	Other details
(Study period)		clusters (%)	
Japan $(2004 - 11)$ [1]	MSM (261)	38.3	Clusters are also linked to China.
, , , , ,	HET (103)	8.7	
Austria $(2008 - 14)$ [2]	MSM (122)	57.4	P < 0.001
	HET (115)	32.2	
Belgium $(2013 - 15)$ [3]	MSM (698)	57.7	$P < 0.001$. Cluster size ≥ 3 .
	HET (508)	22.4	The largest cluster is sized 94,
			consisting of 68 MSM and 15 HET.
Cyprus $(1986 - 2012)$ [4]	MSM (164)	36.6	Cluster size ≥ 3 . The two largest clusters
	HET (129)	2.3	are sized 13, having $11 - 12$ MSM.
France $(1999 - 2014)$ [5]	MSM (636)	32.1	$P < 0.01$. Cluster size ≥ 3 .
	HET (267)	3.8	The largest cluster, sized 41,
			has 39 MSM and a HET.
Germany $(1999 - 2016)$	MSM (1, 448)	27.3	
[6]	HET (622)	18.2	
Netherlands $(1981 - 2011)$	MSM (4, 288)	51.2	Cluster size ≥ 10 .
[7]	HET (849)	55.8	
Spain $(2004 - 14)$ [8]	MSM (637)	_	8 of 12 clusters sized \geq 10 consist only
	HET (204)	_	of MSM. The largest cluster has a size
			of 111, and is composed only of MSM.
USA $(2001 - 15)$ [9]	MSM(1,597)	37.4	The largest cluster has a size
	HET (953)	28.3	of 39, and has only MSM.
Canada and 9	MSM (4, 980)	51.4	The 9 countries are Italy, Greece,
European countries	Bisexual	27.7	Netherlands, Germany, France,
(1981 - 2011) [10]	men $(2,087)$		UK, Norway, Austria, and Spain.
			P < 0.001.

References

- 1. Kondo, M. et al. Emergence in Japan of an HIV-1 variant associated with transmission among men who have sex with men (MSM) in China: first indication of the international dissemination of the Chinese MSM lineage. J. Virol. 87, 5351-5361 (2013). URL https://jvi.asm.org/content/87/10/5351.
- 2. Hoenigl, M. et al. Characterization of HIV transmission in South-east Austria. PLoS ONE 11, e0151478 (2016). URL https://doi.org/10.1371/journal.pone.0151478.
- 3. Verhofstede, C. et al. Phylogenetic analysis of the Belgian HIV-1 epidemic reveals that local transmission is almost exclusively driven by men having sex with men despite presence of large African migrant communities. Infect. Genet. Evol. 61, 36–44 (2018). URL https://doi.org/10.1016/j.meegid.2018.03.002.

- 4. Pineda-Peña, A. et al. HIV-1 infection in Cyprus, the Eastern Mediterranean European frontier: a densely sampled transmission dynamics analysis from 1986 to 2012. Sci. Rep. 8, 1702 (2018). URL https://doi.org/10.1038/s41598-017-19080-5.
- 5. Chaillon, A. et al. Spatiotemporal dynamics of HIV-1 transmission in France (1999-2014) and impact of targeted prevention strategies. Retrovirology 14, 15 (2017). URL https://doi.org/10.1186/s12977-017-0339-4.
- 6. Stecher, M. et al. Molecular epidemiology of the HIV epidemic in three German metropolitan regions Cologne/Bonn, Munich and Hannover, 1999–2016. Sci. Rep. 8, 6799 (2018). URL https://doi.org/10.1038/s41598-018-25004-8.
- 7. Bezemer, D. et al. Dispersion of the HIV-1 epidemic in men who have sex with men in the Netherlands: a combined mathematical model and phylogenetic analysis. PLoS Med. 12, e1001898 (2015). URL https://doi.org/10.1371/journal.pmed.1001898.
- 8. Patino-Galindo, J. A. et al. The molecular epidemiology of HIV-1 in the Comunidad Valenciana (Spain): analysis of transmission clusters. Sci. Rep. 7, 11584 (2017). URL https://doi.org/10.1038/s41598-017-10286-1.
- 9. Dennis, A. M. et al. HIV-1 transmission clustering and phylodynamics highlight the important role of young men who have sex with men. AIDS Res. Hum. Retroviruses 34, 879–88 (2018). URL https://doi.org/10.1089/aid.2018.0039.
- 10. Paraskevis, D. et al. HIV-1 molecular transmission clusters in nine European countries and Canada: association with demographic and clinical factors. BMC Med. 17, 4 (2019). URL https://doi.org/10.1186/s12916-018-1241-1.