

Table S4. Per-site nucleotide divergence (d) and standard deviation (s.d.) among chickens, RJF and GJF.

Intron	Per-site nucleotide divergence (d)					
	Chickens vs. RJF	s.d.	Chickens vs. GJF	s.d.	RJF vs. GJF	s.d.
1	0.008	0.004	0.005	0.004	0.007	0.003
2	0.01	0.004	0.013	0.006	0.015	0.004
3	0.015	0.005	0.015	0.008	0.028	0.009
4	0.003	0.002	N.A. ^a	N.A.	N.A.	N.A.
5	0.003	0.006	0.005	0.006	0.002	0.002
6	0.003	0.002	0.003	0.002	0.003	0.001
7	0.004	0.003	0.009	0.003	0.009	0.003
8	0.011	0.006	0.026	0.004	0.022	0.005
9	0.008	0.004	0.016	0.006	0.017	0.007
10	0.017	0.003	0.025	0.008	0.029	0.008
11	0.007	0.003	0.012	0.004	0.011	0.002
12	0.01	0.005	0.017	0.007	0.02	0.007
13	0.017	0.01	0.024	0.01	0.022	0.009
14	0.006	0.004	0.015	0.006	0.014	0.003
15	0.006	0.004	0.022	0.014	0.022	0.014
16	0.006	0.003	0.012	0.005	0.013	0.005
17	0.015	0.005	0.026	0.009	0.023	0.007
18	0.005	0.003	N.A.	N.A.	N.A.	N.A.
19	0.008	0.005	0.015	0.006	0.014	0.005
20	0.007	0.004	0.018	0.007	0.017	0.008
21	0.005	0.001	0.01	0.005	0.011	0.003
22	0.01	0.004	0.018	0.008	0.019	0.006
23	0.011	0.006	0.02	0.008	0.018	0.005
24	0.004	0.003	0.031	0.015	0.029	0.014
25	0.005	0.002	0.012	0.004	0.012	0.004
26	0.012	0.005	0.03	0.011	0.031	0.011
27	0.002	0.004	0.009	0.003	0.009	0.003
28	0.012	0.006	0.018	0.007	0.011	0.004
29	0.001	0.001	0.004	0.001	0.004	0.001
30	0.004	0.003	0.015	0.005	0.015	0.005

^a N.A. indicates “not available”.